


AMENDMENTS TO THE CLAIMS

This listing of claims replaces all prior listings and versions.

1. (previously presented): A method of reducing liposome-induced complement activation upon in vivo administration of liposomes containing an entrapped chemotherapeutic agent,

comprising providing liposomes comprised of a vesicle-forming lipid and between 1-10 mole percent of a neutral lipopolymer having the formula:



where each of R¹ and R² is an alkyl or alkenyl chain having between 8 and 24 carbon atoms;

n=10-300,

Z is selected from the group consisting of C₁-C₃ alkoxy, C₁-C₃ alkyl ether, n-methylamide, dimethylamide, methylcarbonate, dimethylcarbonate, carbamate, amide, n-methylacetamide, hydroxy, benzyloxy, carboxylic ester, C₁-C₃ alkyl carbonate, and aryl carbonate; and

L is selected from the group consisting of (i) -X-(C=O)-Y-CH₂-, (ii) -X-(C=O)-, and (iii) -X-CH₂-, where X and Y are independently selected from oxygen, NH, and a direct bond, with the proviso that when L is -X-(C=O)-, X is not NH; and the remainder vesicle-forming lipids.

2. (original): The method of claim 1, wherein X is oxygen and Y is nitrogen.
3. (original): The method of claim 1, wherein L is a carbamate linkage, an ester linkage, or a carbonate linkage.
4. (original): The method of claim 3, wherein L is -O-(C=O)-NH-CH_2 (a carbamate linkage).
5. (original): The method of claim 1, wherein Z is hydroxy or methoxy.
6. (currently amended): The method of claim 1, wherein said ~~preparing includes~~ ~~preparing~~ liposomes containing about 1 to 10 mole % of the neutral lipopolymer distearoyl (carbamate-linked) polyethylene glycol.
7. (currently amended): The method of claim 1, wherein said ~~preparing includes~~ ~~preparing~~ liposomes containing about 1 to 10 mole % of the neutral lipopolymer methoxy-polyethelene glycol 1,2 distearoyl glycerol.
8. (original): The method of claim 1, wherein each of R^1 and R^2 is an unbranched alkyl or alkenyl chain having between 8 and 24 carbon atoms.
9. (original): The method of claim 8 wherein each of each of R^1 and R^2 is $\text{C}_{17}\text{H}_{35}$.
10. (original): The method of claim 1, wherein n is between about 20 and about 115.
11. (canceled).
12. (currently amended): The method of claim 1 ~~[[11]]~~, wherein said chemotherapeutic agent is an anthracycline antibiotic.

13. (original): The method of claim 12, wherein said chemotherapeutic agent selected from the group consisting of doxorubicin, daunorubicin, epirubicin, and idarubicin.

14. (currently amended): The method of claim 1 [[11]], wherein said chemotherapeutic agent is a platinum-containing compound.

15. (original): The method of claim 14, wherein said platinum-containing antibiotic is cisplatin or a cisplatin analogue selected from the group consisting of carboplatin, ormaplatin, oxaliplatin, ((-)-(R)-2-aminomethylpyrrolidine (1,1-cyclobutane dicarboxylato))platinum, zeniplatin, enloplatin, lobaplatin, (SP-4-3(R)-1,1-cyclobutane-dicarboxylato(2-)-(2-methyl-1,4-bu- tanediamine-N,N'))platinum, nedaplatin and bis-acetato-ammine-dichloro-cyc- lohexylamine-platinum(IV).

16 to 27. (canceled).